

hMSC Chondrogenesis Differentiation Kit

Product Manual

Catalog#RP02014-B 1 Kit

I. Product Introduction

Shownin hMSC Chondrogenic Differentiation Kit exhibits high-efficiency chondrogenic lineage-specific differentiation capacity, and is applicable for the induction of chondrogenic differentiation in human mesenchymal stem cells (hMSC).

II. Product Information

Table 1: hMSC Chondrogenesis Differentiation Kit Product Description

Product Information	Cat.No.	Amount	Storage
hMSC Chondrogenesis Differentiation Kit contains:	RP02014-B	1 Kit	*
Chondrogenesis Differentiation Basal Medium	RP02014-B-01	80 mL	2 °C to 8 °C
Chondrogenesis Differentiation Supplement	RP02014-B-02	20 mL	-80 °C to -20 °C

*After mixing the basal medium and the supplement to form the complete medium, it can be stored at 2–8 °C and should be used up within 2 weeks.

III. Reagents and Materials

Table 2: Reagents & Materials

Reagents & Materials	Brand (e.g.)	Cat.No. (e.g.)
NcMission™ hMSC Medium V3.0	Shownin	RP02010
Alcian Blue 8GX Solution	OriCell	No.ALCB-10001
4% PFA solution	Biosharp	BL539A
1× DPBS w/o Ca2+ /Mg2+	Thermo Sci.	14190250
6-well Plate	Thermo Sci.	140685
1 mL/5 mL/10 mL/25 mL Pipettes	Thermo Sci.	N/A
15 mL/50 mL Centrifuge Tubes	Thermo Sci.	N/A
10 µL/200 µL/1000 µL Pipette Tips	Rainin.	N/A

IV. Reagent Preparation

(i) Preparation of Complete Medium for hMSC Chondrogenesis Differentiation

1. Thaw the Chondrogenesis Differentiation Supplement at 4 °C. **Do not thaw it at 37 °C.**
2. In a biosafety cabinet, use a sterile pipette to mix the following components to prepare 100 mL of complete differentiation medium.

Chondrogenesis Differentiation Basal Medium: 90 mL

Chondrogenesis Differentiation Supplement: 10 mL

3. The complete medium can be stored at 4 °C and should be used within 2 weeks. Protect from light during storage and use.

Tips: The Supplement can be aliquoted and stored frozen according to actual usage. The total number of freeze-thaw cycles should not exceed 2.

V. Chondrogenesis Differentiation of MSCs

(i) Preparation of MSCs

1. Please refer to the product manual of **NcMission™ hMSC Medium V3.0** in detail.
2. Culture hMSCs in **NcMission™ hMSC Medium V3.0**. Seed hMSCs at a density of 5,000–10,000 cells/cm² in a 6-well plate. Gently rock the plate in a cross pattern three times and place in a 37 °C, 5% CO₂, humidified incubator. Rock the plate again three times and maintain culture.

(ii) Chondrogenesis Differentiation of MSCs

1. Granulation formation: When hMSCs reach approximately 80% confluence:
 - 1) Digest cells in 1 mL/well of 0.5× TryPLE at 37 °C for 3 minutes.
 - 2) Balance the tubes and centrifuge at 250 × g for 5 minutes (acceleration 3, deceleration 7). Aspirate the supernatant and resuspend cells in NcMission hMSC Medium. Count cells.
 - 3) Transfer 2 × 10⁵ cells into a 15 mL centrifuge tube (2 tubes per group). If the suspension volume is less than 1 mL, resuspend cells in NcMission hMSC Medium to 1 mL. Balance and centrifuge at room temperature (300 × g for 5 minutes; acceleration 3, deceleration 7).
 - 4) Aspirate the supernatant, gently tap the tube to disperse the MSC pellet, and resuspend in 1 mL of complete medium per tube. Balance and centrifuge at room temperature (450 × g for 10 minutes).
 - 5) Loosen the cap and incubate at 37 °C, 5% CO₂ for 16 hours (overnight).

2. Induction of Chondrogenesis (Day 1): Replace the NcMission hMSC Medium with the **hMSC Chondrogenic Differentiation Complete Medium**. Carefully aspirate the NcMission hMSC Medium without disturbing the pellets. Slowly add 2 mL of chondrogenic medium per tube along the wall.
3. Maintain culture for 10–20 days. Change **hMSC Chondrogenic Differentiation Complete Medium** every 2 days, 2 mL/tube. Avoid aspirating pellets during the process (14 days of culture is sufficient for rapid cartilage formation).
4. Fixation, dehydration, staining, microphotography:
 - 1) Fixation and dehydration: Aspirate the medium and wash the pellets with 2 mL/tube of DPBS. Aspirate DPBS and fix with 2 mL/tube of 4% PFA at 25 °C for 10 minutes.
 - 2) Frozen Sectioning: Embed pellets in embedding medium and section at 10 µm thickness (embedded samples can be stored at -80 °C for up to 2 months).
 - 3) Washing: Soak slides in DPBS and wash twice on a shaker (10 minutes each with an appropriate speed).
 - 4) Staining: After drying, apply OriCell Alcian Blue 8GX Solution to each section, ensuring full coverage. Place slides in a humidity chamber and stain at 37 °C for 30 minutes.
 - 5) Cleaning and Photography: Rinse slides under slow running water for 3 minutes. Dry and observe under a microscope. Avoid direct water flow on the sample to prevent detachment.